January 20 – February 1, 2018 | Compiled by Greg Smith





With Greg Smith and participants Barbara, Elna, Catherine, Sanford (Sandy), Cathie, Ginger, Daniel, Muriel, George, Giulio, Lou, and Joan

Species List: 113 Total Birds Seen

Anatidae: Ducks, Geese, and Waterfowl (10)

Greylag Goose (Anser anser) Seen early in the trip on our drive to Tiri Tiri Mantangi.

Canada Goose (Branta canadensis) Usually found near most large bodies of freshwater.

Black Swan (*Cygnus atratus*) They were also found on nearly any large body of water, usually in pairs. **Paradise Shelduck** (*Tadorna variegata*) Scattered through both of the main islands, usually in pairs or small groups.

Blue Duck (*Hymenolaimus malacorhynchos*) We were fortunate in viewing at least twelve of these fast water ducks around Tongoriro.

Australian Shoveler (Anas rhynchotis) Best looks were at birds on our drive from Picton to Kaikoura.

Pacific Black Duck (Anas superciliosa) Seen well in the ponds right after our lunch near Waihu.

Mallard (*Anas platyrhynchos*) More common than most of the other small waterfowl, very domesticated acting.

Gray Teal (Anas gracilis) Seen more regularly early on in the trip as opposed to the end.

New Zealand Scaup (Aythya novaeseelandiae) Very common on Great Lake Taupo and Lake Rotorua.

Spheniscidae: *Penguins (1)*

Fiordland Penguin (*Eudyptes pachyrhynchus*) Seen by a few of the group on Monro Beach.

Odontophoridae: New World Quail (1)

California Quail (*Callipepla californica*) Easy to find on the North Island anytime we were near brushy habitat.

Phasianidae: Pheasants, Grouse, and Allies (1)

Wild Turkey (*Meleagris gallopavo*) Seen a few times a day in paddocks that were bordered by trees on the North Island.

Podicipedidae: Grebes (2)

New Zealand Grebe (*Poliocephalus rufopectus*) Dabchicks were seen a few times on the North Island where there was deep water.

January 20 – February 1, 2018 | Compiled by Greg Smith

Great Crested Grebe (*Podiceps cristatus*) Best looks were at two birds on a small lake on our drive from Picton to Kaikoura.

Diomedeidae: Albatrosses (4)

White-capped Albatross (*Thalassarche cauta cauta*) We had three to four on our albatross outing in Kaikoura.

Salvin's Albatross (Thalassarche salvini) The most common albatross out of Kaikoura.

Royal Albatross (Diomedea epomophora) There were at least three of these, the biggest albatross.

Wandering Albatross (*Diomedea exulans*) The ones we got long looks at were older individuals given the amount of white.

Procellariidae: Shearwaters and Petrels (11)

Northern Giant-Petrel (Macronectes halli)

Cook's Petrel (Pterodroma cookii)

Fairy Prion (Pachyptila turtur)

White-chinned Petrel (Procellaria aequinoctialis)

Black Petrel (Procellaria parkinsoni)

Westland Petrel (*Procellaria westlandica*)

Flesh-footed Shearwater (Ardenna carneipes)

Buller's Shearwater (Ardenna bulleri) Seen sporadically on our albatross outing in Kaikoura.

Sooty Shearwater (*Ardenna grisea*) Two were seen on the ferry crossing from the North to the South Island.

Hutton's Shearwater (*Puffinus huttoni*) They nest near Kaikoura and we saw them waiting for darkness in rafts

Fluttering Shearwater (*Puffinus gavial*) Most common in the Marlborough Sounds from the ferry and wildlife boat trip.

Hydrobatidae: Storm-Petrels (1)

White-faced Storm-Petrel (*Pelagodroma marina*) We had two individuals on our albatross trip out of Kaikoura.

Sulidae: Boobies and Gannets (1)

Australasian Gannet (Morus serrator) The colony at Muriwai was certainly a treat.

Phalacrocoracidae: Cormorants and Shags (7)

Little Pied Shag (*Microcarbo melanoleucos*) Sporadic, but usually around smaller bodies of freshwater or rocky coastline.

Great Cormorant (*Phalacrocorax carbo*) Seen on our wildlife ride in the Marlborough Sounds.

Spotted Shag (*Phalacrocorax punctatus*) First seen in the Marlborough Sounds.

Little Black Cormorant (*Phalacrocorax sulcirostris*) Usually on freshwater for this trip, with best looks at Lake Rotorua.

Pied Cormorant (*Phalacrocorax varius*) The more common shag around saltwater.

January 20 – February 1, 2018 | Compiled by Greg Smith

New Zealand King Shag (*Phalacrocorax carunculatus*) The area in the Marlborough Sounds was where we found a small flock.

Otago Shag (*Phalacrocorax chalconotus*) Great looks at the former Stewart Island Shag on the coast below the albatross centre.

Ardeidae: Herons, Egrets, and Bitterns (2)

Great Egret (*Ardea alba modesta*) We only saw two of these birds, most probably because it was nesting season.

White-faced Heron (*Egretta novaehollandiae*) Usually seen on any fair-sized body of water, with best looks on Great Lake Taupo.

Threskiornithidae: Ibises and Spoonbills (1)

Royal Spoonbill (*Platalea regia*) Not encountered too often, and always in small numbers near freshwater lakes.

Rallidae: Rails, Gallinules, and Coots (2)

Australasian Swamphen (*Porphyrio melanotus*) The Pukekhoe was usually grazing in most areas with short grass.

Weka (*Gallirallus australis*) Two were foraging on the beach of our island stop in Marlborough Sounds. **Eurasian Coot** (*Fulica atra*) Only a few were seen and those were on Great Lake Taupo.

Accipitridae: Hawks, Eagles, and Kites (1)

Swamp Harrier (*Circus approximans*) Seen every day in most non-forest habitats.

Recurvirostridae: Stilts and Avocets (2)

Pied Stilt (*Himantopus leucocephalus*) Large groups of this "chatty" shorebird were seen at the Miranda chernier area.

Black Stilt (*Himantopus novaezelandiae*) A few adults were foraging in the delta of the river that drained the Aoraki glaciers.

Haematopodidae: Oystercatchers (2)

South Island Oystercatcher (*Haematopus finschi*) A large mixed flock at Miranda allowed us to compare with the Variable.

Variable Oystercatcher (*Haematopus unicolor*) The more common of the oystercatchers, especially on the North Island.

Charadriidae: Plovers and Lapwings (5)

Pacific Golden-Plover (*Pluvialis fulva*) Seen at both Miranda and along the edge of the Foxton estuary. **Masked Lapwing** (*Vanellus miles novaehollandiae*) Most large paddocks had a pair, especially on the North Island.

Red-breasted Dotterel (*Charadrius obscurus*) Good looks at both Miranda and along the edge of the estuary near Foxton.

January 20 – February 1, 2018 | Compiled by Greg Smith

Double-banded Plover (Charadrius bicinctus) A good-sized flock at Miranda.

Wrybill (*Anarhynchus frontalis*) This braided river nester was seen well at Miranda in medium-sized flocks.

Scolopacidae: Sandpipers and Allies (5)

Bar-tailed Godwit (*Limosa lapponica*) Always a treat to see these long-distance migrants with a large flock out in Miranda.

Ruddy Turnstone (*Arenaria interpres*) Only seen on the chernier flats at Miranda.

Sanderling (Calidris alba) Three individuals were foraging together in the Foxton Estuary.

Red Knot (Calidris canutus) Seen in basic plumage at Miranda and foraging at the estuary near Foxton.

Sharp-tailed Sandpiper (*Calidris acuminata*) Two, maybe three were seen only at Miranda.

Stercorariidae: Skuas and Jaegers (3)

Brown Skua (*Stercorarius antarcticus lonnbergi*) Seen briefly on our boat crossing to Tiri Tiri Mantangi. **Arctic Skua** (*Stercorarius parasiticus*) A single individual was viewed on our crossing to Tiri Tiri Mantangi. **Parasitic Skua** (*Stercorarius parasiticus*) Seen most every time we were out on saltwater, always in very small numbers.

Laridae: Gulls, Terns, and Skimmers (7)

Black-billed Gull (*Chroicocephalus bulleri*) We only saw this braided river-nester once, on the chenier flats at Miranda.

Red-billed Gull (*Chroicocephalus scopulinus*) Common anywhere around the coast of the New Zealand. **Kelp Gull** (*Larus dominicanus*) Or Southern Black-backed Gull was common on most of the trip when near the coast.

Caspian Tern (Hydroprogne caspia) Only one, and that was at the Miranda Shorebird Centre.

Whiskered Tern (Chlidonias hybrid) A vagrant that had been reported was seen by all near Waihu on Great Lake Taupo.

Black-fronted Tern (*Chlidonias albostriatus*) Not common on this trip with just a few seen near Aoraki. White-fronted Tern (*Sterna striata*) The common tern on the coast.

Columbidae: Pigeons and Doves (3)

Rock Pigeon (*Columba livia*) Cities, parks, and barns.

Spotted Dove (*Streptopelia chinensis*) Only found once and that was as we drove to Muriwai from Tiri Mantangi.

New Zealand Pigeon (*Hemiphaga novaeseelandiae*) If there was native forest, we would always see some of these huge pigeons.

Cuculidae: Cuckoos (1)

Long-tailed Koel (Urodynamis taitensis) Heard in the Pureora Forest on our morning hike.

Strigidae: Owls (1)

January 20 – February 1, 2018 | Compiled by Greg Smith

Southern Boobook (*Ninox novaeseelandiae novaeseelandiae*) Seen by most on Tiri Tiri Mantangi and heard both nights at Lake Moeraki.

Alcedinidae: Kingfishers (1)

Sacred Kingfisher (Todiramphus sanctus) Not at all common, but seen well by all of us.

Falconidae: Falcons and Caracaras (1)

New Zealand Falcon (Falco novaeseelandiae) Heard calling repeatedly on the hike to Monro Beach.

Strigopidae: New Zealand Parrots (2)

Kea (*Nestor notabilis*) During our lunch stop in Arthur's Pass, we got to see one individual working under some shrubs.

New Zealand Kaka (*Nestor meridionalis*) The Pureora Forest had a few of these parrots perched atop of the matawi trees.

Psittaculidae: Old World Parrots (4)

Orange-fronted Parakeet (*Eupsittula canicularis*) All saw one of the 300+ left in the world on an island in Marlborough Sounds.

Red-crowned Parakeet (*Cyanoramphus novaezelandiae novaezelandiae*) Seen crawling around the trees and calling at Pureora.

Yellow-crowned Parakeet (*Cyanoramphus auriceps*) During our breakfast in Pureora Forest we got first of three looks at this species.

Acanthisittidae: New Zealand Wrens (1)

Rifleman (Acanthisitta chloris) Seen on Tiri Tiri Mantangi.

Meliphagidae: Honeyeaters (2)

Tui (*Prosthemadera novaeseelandiae*) Best looks were certainly on Tiri Tiri Mantangi, but we did see them in other locales.

New Zealand Bellbird (*Anthornis melanura*) The feeding stations on Tiri Tiri Mantangi gave us some up close looks.

Acanthizidae: Thornbills and Allies (1)

Greygone (Gerygone igata) Seen by the group on the trail near Blue Pools on the drive to Lake Ohau.

Mohouidae: Whiteheads (3)

Whitehead (Mohoua albicilla) Family groups seemed to be everywhere on our Tiri Tiri Mantangi visit. Pipipi (Mohoua novaeseelandiae) Seen by the group on the trail near Blue Pools on the drive to Lake Ohau.

Callaeidae: Wattlebirds (3)

January 20 – February 1, 2018 | Compiled by Greg Smith

North Island Kokako (*Callaeas wilsoni*) Really good looks at a family group feeding on the fruits of the cabbage tree.

North Island Saddleback (*Philesturnus rufusater*) Great looks while out on Tiri Tiri Mantangi **South Island Saddleback** (*Philesturnus carunculatus*) We got to see this South Island relative down on some of the predator-free islands.

Notiomystidae: Stitchbird (1)

Stitchbird (*Notiomystis cincta*) Only on Tiri Tiri Mantangi.

Cracticidae: Bellmagpies and Allies (1)

Australian Magpie (*Gymnorhina tibicen*) Most paddocks on both islands had a resident pair of these birds.

Rhipiduridae: Fantails (1)

New Zealand Fantail (*Rhipidura fuliginosa*) Not common, but seen every time we were in native forest.

Petroicidae: Australasian Robins (2)

Tomtit (*Petroica macrocephala*) Seen in the Marlborough Sounds and at Lake Moeraki.

New Zealand Robin (Petroica australis) First seen on Tiri Tiri Mantangi.

Alaudidae: Larks (1)

Skylark (*Alauda arvensis*) Seen during a coffee/tea stop on the drive from Kaikoura.

Hirundinidae: Swallows (1)

Welcome Swallow (Hirundo neoxena) They were everywhere there was a body of water nearby.

Locustellidae: Grassbirds and Allies (1)

Fernbird (*Megalurus punctatus*) Only a pair was seen, and that was on our morning breakfast stop in Pureora Forest.

Zosteropidae: White-eyes, Yuhinas, and Allies (1)

Silver-eye (*Zosterops lateralis*) When seen, we usually saw them in family groups or flocks.

Turdidae: Thrushes and Allies (2)

Eurasian Blackbird (*Turdus merula*) A front yard bird throughout most of New Zealand. **Song Thrush** (*Turdus philomelos*) Another introduced species, but not nearly as common as the blackbird.

Sturnidae: Starlings (2)

European Starling (*Sturnus vulgaris*) Most everywhere there were fields and/or hedgerows. **Common Myna** (*Acridotheres tristis*) Haven't seen any on the South Island, but they were very common in Auckland.

January 20 – February 1, 2018 | Compiled by Greg Smith

Motacillidae: Wagtails and Pipits (1)

Australasian Pipit (Anthus novaeseelandiae) Seen while looking at a couple of Blue Duck in Tongariro.

Emberizidae: Buntings and New World Sparrows (1)

Yellowhammer (Emberiza citrinella) Seen sporadically feeding alongside fallow paddocks.

Fringillidae: Finches, Euphonias, and Allies (4)

Common Chaffinch (*Fringilla coelebs*) Seen by the group on the trail near Blue Pools on the drive to Lake Ohau.

European Greenfinch (*Chloris chloris*) Seen only a couple of times, and only on the North Island.

Common Redpoll (Acanthis flammea) Seen during our lunch stop in Wairu.

European Goldfinch (*Carduelis carduelis*) Flocks were doing a pretty good job on grass-covered paddocks.

Passeridae: Old World Sparrows (1)

House Sparrow (Passer domesticus) Unfortunately, everywhere. ...